

AMENDMENTS TO THE CLAIMS

Please amend the claims as follows:

LISTING OF CLAIMS:

1. (Currently Amended): A process for the production of L-ascorbic acid comprising:

(a) contacting an enzyme with a substrate which is selected from the group consisting of L-gulose, L-galactose, L-idose, and L-talose;

(b) converting the substrate directly into L-ascorbic acid by catalytic ~~catalytic~~ activity of the enzyme under suitable culture conditions; and

(c) isolating L-ascorbic acid from the reaction mixture,

wherein said enzyme has (1) the amino acid sequence of SEQ ID NO: 2 or (2) an amino acid sequence with 90% sequence identity to SEQ ID NO: 2 and with the activity to produce L-ascorbic acid or (3) an amino acid sequence encoded by the DNA sequence of SEQ ID NO: 1 or (4) an amino acid sequence encoded by a DNA sequence that hybridizes under stringent hybridization and wash conditions to the DNA sequence of SEQ ID: 1 and having the activity to produce L-ascorbic acid.

2. (Currently Amended): A process for the production of L-ascorbic acid with an enzyme having (1) the amino acid sequence of SEQ ID NO: 2 or (2) an amino acid sequence with 90% sequence identity to SEQ ID NO: 2 and with the activity to produce L-ascorbic acid or (3) an amino acid sequence encoded by the DNA sequence of SEQ ID NO: 1 or (4) an amino acid sequence encoded by a DNA sequence that hybridizes under stringent hybridization and wash conditions to the DNA sequence of

SEQ ID: 1 and having the activity to produce L-ascorbic acid, whereby L-ascorbic acid is produced from a substrate which is selected from the group consisting of L-gulono-1,4-lactone, L-gulonic acid, L-galactono-1,4-lactone, L-galactonic acid, L-idono-1,4-lactone, L-idonic acid, L-talono-1,4-lactone, and L-talonic acid,

said process comprising the steps of:

- (a) contacting the enzyme with the substrate,
- (b) converting the substrate directly into L-ascorbic acid by catalytic ~~catalytical~~ activity of the enzyme under suitable culture conditions; and
- (c) isolating L-ascorbic acid from the reaction mixture.

3. (Withdrawn): A process for the production of L-gulono-1,4-lactone or L-gulonic acid with an enzyme having the amino acid sequence of SEQ ID NO: 2 or an amino acid sequence that is 90% identical thereto, with the activity to produce L-gulono-1,4-lactone or L-gulonic acid, whereby L-gulono-1,4-lactone or L-gulonic acid is produced from L-gulose.

4. (Withdrawn): A process for the production of L-galactono-1,4-lactone or L-galactonic acid with an enzyme having the amino acid sequence of SEQ ID NO: 2 or an amino acid sequence that is 90% identical thereto, with the activity to produce L-galactono-1,4-lactone or L-galactonic acid, whereby L-galactono-1,4-lactone or L-galactonic acid is produced from L-galactose.

5. (Cancelled).

6. (Previously presented): A process according to claim 1, wherein the process is conducted for 1 to 120 h at a pH of about 1 to about 9 and at a temperature of about 13°C to about 45°C.

7. (Original): A process according to claim 6, wherein the process is conducted at a pH of about 2 to about 8 and at a temperature of about 18°C to about 42°C.

8. (Currently amended): A process for producing L-ascorbic acid comprising (a) contacting a substrate which is selected from the group consisting of L-gulose, L-galactose, L-idose, L-talose, L-gulono-1,4-lactone, L-gulonic acid, L-galactono-1,4-lactone, and L-galactonic acid with an enzyme derivable from *G. oxydans* DSM 4025, (b) converting the substrate directly into L-ascorbic acid by catalytic ~~catalytic~~ activity of the enzyme under suitable culture conditions and (c) isolating L-ascorbic acid from the reaction mixture; wherein the enzyme has the following physico-chemical properties:

- (a) molecular weight of about 60,000 Da on SDS-PAGE;
- (b) substrate specificity for primary and secondary alcohols and aldehydes;
- (c) pH-stability at pH of about 6 to about 9;
- (d) pH-optimum at pH of about 8.0; and
- (e) inhibited by Cu^{2+} , Zn^{2+} , Mn^{2+} , Fe^{2+} , and Fe^{3+} .

9. (Withdrawn): A process for producing L-gulono-1,4-lactone or L-gulonic acid comprising contacting L-gulose with Enzyme B of *G. oxydans* DSM 4025 and isolating L-gulono-1,4-lactone or L-gulonic acid from the reaction mixture, wherein Enzyme B has the following physico-chemical properties:

- (a) molecular weight of about 60,000 Da on SDS-PAGE;

(b) substrate specificity for primary and secondary alcohols and aldehydes;

(c) pH-stability at pH of about 6 to about 9;

(d) pH-optimum at pH of about 8.0; and

(e) inhibited by Cu^{2+} , Zn^{2+} , Mn^{2+} , Fe^{2+} , and Fe^{3+} .

10. (Withdrawn): A process for producing L-galactono-1,4-lactone or galactonic acid comprising contacting L-galactose with Enzyme B of *G. oxydans* DSM 4025 and isolating L-galactono-1,4-lactone or galactonic acid from the reaction mixture, wherein Enzyme B has the following physico-chemical properties:

(a) molecular weight of about 60,000 Da on SDS-PAGE;

(b) substrate specificity for primary and secondary alcohols and aldehydes;

(c) pH-stability at pH of about 6 to about 9;

(d) pH-optimum at pH of about 8.0; and

(e) inhibited by Cu^{2+} , Zn^{2+} , Mn^{2+} , Fe^{2+} , and Fe^{3+} .

11. (Withdrawn): A process according to claim 3 comprising (a) contacting the enzyme with the substrate and (b) isolating the L-gulono-1,4-lactone or L-gulonic acid from the reaction mixture.

12. (Withdrawn): A process according to claim 4 comprising (a) contacting the enzyme with the substrate and (b) isolating the L-galactono-1,4-lactone or L-galactonic acid from the reaction mixture.

13. (Previously presented): A process according to claim 2, wherein the process is conducted for 1 to 120 h at a pH of about 1 to about 9 and at a temperature of about 13°C to about 45°C.

14. (Withdrawn): A process according to claim 3, wherein the process is conducted for 1 to 120 h at a pH of about 1 to about 9 and at a temperature of about 13°C to about 45°C.

15. (Withdrawn): A process according to claim 4, wherein the process is conducted for 1 to 120 h at a pH of about 1 to about 9 and at a temperature of about 13°C to about 45°C.

16. (Previously presented): A process according to claim 8, wherein the process is conducted for 1 to 120 h at a pH of about 1 to about 9 and at a temperature of about 13°C to about 45°C.

17. (Withdrawn): A process according to claim 11, wherein the process is conducted for 1 to 120 h at a pH of about 1 to about 9 and at a temperature of about 13°C to about 45°C.

18. (Withdrawn): A process according to claim 12, wherein the process is conducted for 1 to 120 h at a pH of about 1 to about 9 and at a temperature of about 13°C to about 45°C.